arXiv:1209.1612v1 [math.RT] 7 Sep 2012

ON THE n-TH DIMENSIONAL POROUS MEDIUM DIFFUSION EQUATION AND GLOBAL ACTIONS OF THE SYMMETRY GROUP

JOSE A. FRANCO

Abstract. The Lie point symmetries of the n-th dimensional porous medium diffusion equation are calculated. By restricting to a special class of smooth functions, the local action of the symmetry group is globalized. This special class of functions is constructed using parabolic induction.

1. Introduction

The theory of Lie groups finds its genesis when Sophus Lie sets up the task to develop an analogue of Galois Theory for differential equations. The original prolongation algorithm of Sophus Lie provides a set of point symmetries of a differential equation (for a modern treatment, see [6]). It is well-known that these point symmetries generate a *local* Lie group action on the space of solutions of the differential equation and this action seldom globalizes. However, most of the modern results on Lie group theory apply to *global* Lie groups and *global* Lie group representations. As a result, many of the standard techniques of representation theory are not always applicable to differential equations. Therefore, it is an important problem to find a globalization of the local action of the symmetry group of a differential equation. This is usually achieved by restricting to a special class of functions.

In this paper, we study the globalization problem for the n-th dimensional porous medium equation,

$$u_t = \Delta_n(u^m), \tag{1.1}$$

where Δ_n is the n-th dimensional Laplacian and u is a function of $x = (x_1, ..., x_n) \in \mathbb{R}^n$ and $t \in \mathbb{R}$. When $m > 1$, this equation models slow diffusion phenomena so this condition is often assumed (c.f. [9]). However, for reasons that will become evident below, we will allow $m \in \mathbb{R} \setminus \{0, 1\}$. In fact, we will be particularly interested in the case $m = \frac{n-2}{n+2}$, because for this special value of m the symmetry group is n-dimensions larger than the generic case. The goal of this paper is to describe the class of functions on which the action of the symmetry group globalizes and to describe the action of the group on these functions. This idea is quite profitable when applied to linear PDE's (see [2],[4], and [8]). However, it has also been used with non-linear PDE's (see [7]).

The paper is organized as follows. In Section 2, we list the infinitesimal generators of the symmetry group of Equation (1.1). For the one-dimensional case, this

---

1991 *Mathematics Subject Classification.* 22E70, 35Q35 .

Key words and phrases. Globalizations, Porous medium equation, Lie group, Lorentz group, parabolic induction.

is well-known (see [1] or [3]). However, we were unable to find a reference for the classification of the symmetry group for $n \geq 2$, therefore, calculate it explicitly here. Then, we embed the symmetry group as a subgroup of $G := SL(2,\mathbb{R}) \times SO(n+1,1)_0$.

In Section 3, we briefly study the structure of the group G. In Section 4, we use parabolic induction to construct a family of representations of G and by restriction, of the symmetry group of (1.1). We use these representations to define class a of smooth functions on which the action of the symmetry group globalizes (see Equation (4.1)). Finally, in Section 5, we write explicitly the action of the 1-parameter subgroups generated by the infinitesimal generators of G.

2. Symmetry Group

Using Lie's prolongation algorithm, we calculate the symmetry group of Equation (1.1); this yields two cases depending on the value of m. When $m \neq \frac{n-2}{n+2}$, the infinitesimal generators of the symmetry group are

$$X_1 = \partial_t \tag{2.1}$$

$$X_2 = E_n + \frac{2u}{m-1}\partial_u \tag{2.2}$$

$$X_3 = -t\partial_t + \frac{u}{m-1}\partial_u \tag{2.3}$$

$$Y_i = \partial_i \text{ for } 1 \leq i \leq n \tag{2.4}$$

$$Z_{i,j} = x_i\partial_j - x_j\partial_i \text{ for } 1 \leq i < j \leq n \tag{2.5}$$

where E_n denotes the n-th dimensional Euler operator. In this case, they span an algebra isomorphic to a parabolic subalgebra of $\mathfrak{g} := \mathfrak{s} \oplus \mathfrak{so}(n+1,1)$, where $\mathfrak{s}$ is isomorphic to a parabolic subalgebra of $\mathfrak{sl}_2(\mathbb{R})$.

When $m = \frac{n-2}{n+2}$, the symmetry group is generated by the differential operators in Equations (2.1) to (2.5), together with:

$$W_i = (x_i^2 - \sum_{j\neq i} x_j^2)\partial_i + \sum_{j\neq i}[2x_ix_j\partial_j] + \frac{4x_iu}{m-1}\partial_u \tag{2.6}$$

for $1 \leq i \leq n$. These infinitesimal generators span an algebra isomorphic to $\mathfrak{g}$. The isomorphism for the $\mathfrak{so}(n+1,1)$ part is explicitly defined in the following way:

$$X_2 \mapsto -E_{n+1,n+2} - E_{n+2,n+1} \tag{2.7}$$

$$\frac{1}{2}(W_i + Y_i) \mapsto E_{n+1,i} - E_{i,n+1} \text{ for } 1 \leq i \leq n \tag{2.8}$$

$$\frac{1}{2}(W_i - Y_i) \mapsto E_{n+2,i} + E_{i,n+2} \text{ for } 1 \leq i \leq n \tag{2.9}$$

$$Z_{i,j} \mapsto E_{i,j} - E_{j,i} \text{ for } 1 \leq i < j \leq n, \tag{2.10}$$

where $E_{k,l}$ is the $(n+2)\times(n+2)$ matrix with single non-zero entry in the k-th row and l-th column. There are two possible choices for $\mathfrak{s}$; we embed it as the upper triangular Borel subalgebra of $\mathfrak{sl}_2(\mathbb{R})$.

3. The Group

With an eye toward the construction of a family of induced representations of the group $G := SL(2,\mathbb{R}) \times SO(n+1,1)_0$, we will study its structure in this section. Since most of this material is standard, most details will be omitted (for a general treatment see VI of [5]).

Let $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ be the Cartan decomposition under the standard Cartan involution. Then, $\mathfrak{k}$ is isomorphic to $\mathfrak{so}(2) \times \mathfrak{so}(n+1)$ with $\mathfrak{so}(2) \subset \mathfrak{sl}_2(\mathbb{R})$ and $\mathfrak{so}(n+1)$ embedded in $\mathfrak{so}(n+1,1)$ in the upper left $(n+1) \times (n+1)$ block. The maximal abelian subalgebra $\mathfrak{a} \subset \mathfrak{p}$ is given by

$$\mathfrak{a} := \left\{ H_{a,s} := \left[\begin{pmatrix} a & 0 \\ 0 & -a \end{pmatrix}, \left(\begin{array}{cccc|c} & & & & 0 \\ & & & & \vdots \\ & 0_{(n+1)\times(n+1)} & & & 0 \\ & & & & -s \\ \hline 0 & \cdots & 0 & -s & 0 \end{array} \right) \right] \mid a, s \in \mathbb{R} \right\}.$$

The centralizer of $\mathfrak{a}$ in $\mathfrak{k}$, $\mathfrak{m}$, is trivial in the $\mathfrak{sl}_2(\mathbb{R})$-component; in the $\mathfrak{so}(n+1,1)$-component, it is isomorphic to $\mathfrak{so}(n)$ and it is embedded in the upper left $n \times n$ block.

If $\nu_i^{\pm} := E_{n+1,i} - E_{i,n+1} \pm E_{n+2,i} \pm E_{i,n+2} \in \mathrm{Mat}_{(n+2)\times(n+2)}$ for $1 \leq i \leq n$, then,

$$\mathfrak{n}^{+} = \left\{ \eta_i^{+} := \left[\begin{pmatrix} 0 & s \\ 0 & 0 \end{pmatrix}, a\nu_i^{+} \right] \mid a, s \in \mathbb{R}, 1 \leq i \leq n \right\}$$

and

$$\mathfrak{n}^{-} = \left\{ \eta_i^{-} := \left[\begin{pmatrix} 0 & 0 \\ s & 0 \end{pmatrix}, a\nu_i^{-} \right] \mid a, s \in \mathbb{R}, 1 \leq i \leq n \right\}.$$

Consequently, the minimal parabolic subalgebras of $\mathfrak{g}$ are defined as $\mathfrak{q} = \mathfrak{m} \oplus \mathfrak{a} \oplus \mathfrak{n}$ and $\mathfrak{q}^{-} = \mathfrak{m} \oplus \mathfrak{a} \oplus \mathfrak{n}^{-}$.

At the level of the group, these subalgebras exponentiate to:

$$A := \left\{ h_{a,y} := \left[\begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix}, \left(\begin{array}{c|cc} I_n & \multicolumn{2}{c}{0} \\ \hline 0 & \cosh y & \sinh y \\ & \sinh y & \cosh y \end{array} \right) \right] \mid y \in \mathbb{R}, a > 0 \right\},$$

$$M := \left\{ m_{j,B} := \left[\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}^{j}, \left(\begin{array}{c|c} B & 0 \\ \hline 0 & I_2 \end{array} \right) \right] \mid B \in SO(n), j \in \mathbb{Z}_2 \right\},$$

$$N := \left\{ n_{t,x} := \left[\begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix}, \left(\begin{array}{c|cc} I_n & -x & x \\ \hline x^T & 1 - \frac{1}{2}\|x\|^2 & \frac{1}{2}\|x\|^2 \\ x^T & -\frac{1}{2}\|x\|^2 & 1 + \frac{1}{2}\|x\|^2 \end{array} \right) \right] \mid (t,x) \in \mathbb{R}^{1,n} \right\},$$

and

$$N^{-} := \left\{ n_{t,x}^{-} := \left[\begin{pmatrix} 1 & 0 \\ t & 1 \end{pmatrix}, \left(\begin{array}{c|cc} I_n & x & x \\ \hline -x^T & 1 - \frac{1}{2}\|x\|^2 & -\frac{1}{2}\|x\|^2 \\ x^T & \frac{1}{2}\|x\|^2 & 1 + \frac{1}{2}\|x\|^2 \end{array} \right) \right] \mid (t,x) \in \mathbb{R}^{1,n} \right\}.$$

The minimal parabolic subgroups corresponding to $\mathfrak{q}$ and $\mathfrak{q}^{-}$ are $Q = MAN$ and $Q^{-} = MAN^{-}$ respectively.

4. Induced Representations

In this section we will construct a family of characters on Q^-. Then, we will induce a representation of G from each of these characters. Via intertwining isomorphisms, we will identify each of these induced representations with a subspace of $C^\infty(\mathbb{R}^{1,n})$. From the standard representation theoretical results (c.f. [5]), we know that G acts globally on these special classes of functions. Since, the symmetry group of the porous medium equation can be embedded in G, the action of G descends to a global action of the symmetry group of the porous medium equation.

We start by noticing that a general element $\varphi \in \mathfrak{a}^*$ is a linear functional that acts by $\varphi(H_{v,y}) = bv + cy$ for fixed constants $b, c \in \mathbb{C}$. By exponentiating $\varphi_{b,c}$ we obtain a character on A. The resulting character, $\chi_{r,s} : A \to \mathbb{C}$, is determined by two continuous parameters $r, s \in \mathbb{C}$ and defined by

$$\chi_{r,s}(h_{a,y}) = a^r e^{sy}.$$

A character $\chi_p : M \to \mathbb{C}$ can be defined by

$$\chi_p(m_{j,B}) = (-1)^{jp},$$

for $p \in \mathbb{Z}_2$. Therefore, we will consider the characters parametrized by two continuous parameters $r, s \in \mathbb{C}$ and a discrete parameter $p \in \mathbb{Z}_2$, $\chi_{p,r,s} : Q^- \to \mathbb{C}$ defined by

$$\chi_{p,r,s}(q^-) = \chi_p(m)\chi_{r,s}(a)$$

where $q^- = man^-$.

We consider the induced representation

$$\begin{aligned}\operatorname{Ind}_{Q^-}^G(\chi_{p,r,s}) := \{\varphi \in C^\infty(G) \mid \varphi(gq^-) = \chi_{p,r,s}(q^-)^{-1}\varphi(g) \\ \text{for } g \in G \text{ and } q^- \in Q^-\},\end{aligned}$$

with the G-action defined by left translation.

The unipotent group N is isomorphic to $\mathbb{R}^{1,n}$ via $(t,x) \mapsto n_{t,x}$. Since Q^-N embeds in G as an open dense set, it is easy to see from the definition of $\operatorname{Ind}_{Q^-}^G(\chi_{p,r,s})$ that an element $\varphi \in \operatorname{Ind}_{Q^-}^G(\chi_{p,r,s})$ is completely determined by its value on N. Therefore, we can define an isomorphism of vector spaces between $\operatorname{Ind}_{Q^-}^G(\chi_{p,r,s})$ and the space

$$I'(p,r,s) := \left\{ f \in C^\infty(\mathbb{R}^{1,n}) \mid f(t,x) = \varphi(n_{t,x}) \text{ for some } \varphi \in \operatorname{Ind}_{Q^-}^G(\chi_{p,r,s}) \right\}. \tag{4.1}$$

A G-module structure can be given to $I'(p,r,s)$ so that the map $\varphi \to f$ is intertwining. Therefore, $\operatorname{Ind}_{Q^-}^G(\chi_{p,r,s})$ is isomorphic to $I'(p,r,s)$ as G-module.

5. Actions on $I'(p,r,s)$

In this section, we will describe the actions of G and of $\mathfrak{g}$ on $I'(p,r,s)$. As a result, we will find special values of r and s that will determine the class of functions on which the action of the symmetry group of Equation (1.1), globalizes.

5.1. Actions of $\mathfrak{sl}(2,\mathbb{R})$ and $SL(2,\mathbb{R})$.

In this section, when we consider the action of an element $g \in SL(2,\mathbb{R})$, it should be understood as the action of its image under the natural inclusion map $SL(2,\mathbb{R}) \hookrightarrow G$. The analogue will be true for elements in the Lie algebra. If possible, we will avoid writing this image explicitly to avoid cumbersome notation.

Proposition 1. *An element* $\left(\begin{smallmatrix} a & b \\ 0 & a^{-1} \end{smallmatrix}\right) \in SL(2,\mathbb{R})$ *acts on* $I'(p,r,s)$ *by*

$$\left(\begin{smallmatrix} a & b \\ 0 & a^{-1} \end{smallmatrix}\right).f(t,x) = \operatorname{sgn}(a)^p |a|^r f\left(\frac{t-ab}{a^2}, x\right).$$

Let $\{H,E,F\}$ *be the standard basis for* $\mathfrak{sl}(2,\mathbb{R})$. *Then,* H *acts by the differential operator* $-2t\partial_t + ru\partial_u$ *and* E *acts by* ∂_t *on* $I'(p,r,s)$.

Proof. First, we write $\left(\begin{smallmatrix} a & b \\ 0 & a^{-1} \end{smallmatrix}\right) = \left(\begin{smallmatrix} 1 & ba \\ 0 & 1 \end{smallmatrix}\right)\left(\begin{smallmatrix} a & 0 \\ 0 & a^{-1} \end{smallmatrix}\right)$ and we calculate each action of each element individually. Let $f \in I'(p,r,s)$. Then, using the definition of $I'(p,r,s)$ and the G action on it, we have

$$\left(\begin{smallmatrix} a & 0 \\ 0 & a^{-1} \end{smallmatrix}\right).f(t,x) = \left(\begin{smallmatrix} a & 0 \\ 0 & a^{-1} \end{smallmatrix}\right).\varphi(n_{t,x}) = \varphi(\left[\left(\begin{smallmatrix} a^{-1} & 0 \\ 0 & a \end{smallmatrix}\right), I_{n+2}\right] n_{t,x}).$$

Now we notice that

$$\left[\left(\begin{smallmatrix} a^{-1} & 0 \\ 0 & a \end{smallmatrix}\right), I_{n+2}\right] n_{t,x} = n_{a^{-2}t,x} \cdot \left[((\left(\begin{smallmatrix} a^{-1} & 0 \\ 0 & a \end{smallmatrix}\right), I_{n+2}\right].$$

Using the character $\chi_{p,r,s}$ and the definition of $\operatorname{Ind}_{Q^-}^{G}(\chi_{p,r,s})$, we obtain

$$\left(\begin{smallmatrix} a & 0 \\ 0 & a^{-1} \end{smallmatrix}\right).f(t,x) = \chi_{p,r,s}\left(\left[(\operatorname{sgn}(a)\begin{pmatrix} |a|^{-1} & 0 \\ 0 & |a| \end{pmatrix}, I_{n+2}\right]\right)^{-1} \varphi(n_{a^{-2}t,x}) = \operatorname{sgn}(a)^p |a|^r f(a^{-2}t,x).$$

The calculation for the other element is similar, thus omitted. The result on the Lie algebra action follows by differentiation. □

Remark 1. *Setting* $r = \frac{2}{m-1}$ *we recover the actions of the elements* X_1 *and* X_3 *in the symmetry group. We did not calculate the action of* F *because the space of solutions to the porous medium equation is not invariant under the subalgebra generated by this element. So, for the goals of this paper, the action of this element is not relevant.*

5.2. Actions of $\mathfrak{so}(n+1,1)$ and $SO(n+1,1)_0$.

As in the previous section, in this section when we consider the action of an element $g \in SO(n+1,1)_0$, it should be understood as the action of its image under the natural inclusion map $SO(n+1,1)_0 \hookrightarrow G$. The same convention will be used for the actions of the Lie algebra.

5.2.1. *Actions of* $\mathfrak{m}$ *and* M.

Lemma 1. *Let* $g_{i,j,\theta} = \exp_{SO(n+1,1)}(\theta(E_{i,j} - E_{j,i}))$ *for* $1 \le i < j \le n$. *Then,*

$$[I_2, g_{i,j,-\theta}] \cdot n_{t,x} = n_{t,g_{i,j,-\theta}x} \cdot [I_2, g_{i,j,-\theta}].$$

Proof. This is a straightforward matrix calculation. □

Proposition 2. *An element* $g_{i,j,\theta}$ *for* $1 \le i < j \le n$ *acts on a function* $f \in I'(p,r,s)$ *by*

$$g_{i,j,\theta}.f(t,x) = f(t, g_{i,j,-\theta}x).$$

The element $E_{i,j} - E_{j,i} \in \mathfrak{so}(n+1,1)$ *for* $1 \le i < j \le n$ *acts on* $I'(p,r,s)$ *by the differential operator*

$$x_i\partial_j - x_j\partial_i.$$

Proof. By the definition of $I'(p,r,s)$ and by the action of G on $\operatorname{Ind}_{Q^-}^{G}(\chi_{p,r,s})$ we have

$$g_{i,j,\theta}.f(t,x) = g_{i,j,\theta}.\varphi(n_{t,x}) = \varphi([I_2, g_{i,j,-\theta}].n_{t,x}).$$

The result now follows from the lemma and the definition of $\chi_{p,r,s}$. The result for the Lie algebra follows by taking $\frac{d}{d\theta}|_{\theta=0}$. □

5.2.2. *Actions of $\mathfrak{n}$ and N.* A straightforward matrix calculation shows that

$$n_{t',x_ie_i}n_{t,x} = n_{t-t',x-x_ie_i}$$

where e_i is the standard basis element of $\mathbb{R}^n$. We use this fact to prove the following proposition.

Proposition 3. *The action of $n_{t',x'}$ on $f \in I'(p,r,s)$ is given by*

$$n_{t',x'}.f(t,x) = f(t-t', x-x')$$

and ν_i^+ acts on $I'(p,r,s)$ by the differential operator ∂_i, for $1 \le i \le n$.

5.2.3. *Actions of $\mathfrak{a}$ and A.* Using the fact that

$$h_{1,\epsilon}^{-1}n_{t,x} = n_{t,e^\epsilon x}h_{1,\epsilon}^{-1},$$

we prove the following proposition.

Proposition 4. *The action of $h_{1,\epsilon}$ on $f \in I'(p,r,s)$ is given by*

$$h_{1,\epsilon}.f(t,x) = e^{s\epsilon}f\left(t, e^\epsilon x\right)$$

and $H_{0,1}$ acts on $I'(p,r,s)$ by the differential operator $E_n + su\partial_u$.

Remark 2. *When $s = \frac{2}{m-1}$, this action corresponds to the action of the infinitesimal generator X_2 of the symmetry group of the porous medium equation* (1.1).

5.2.4. *Actions of $\mathfrak{n}^-$ and N^-.* To describe this action, we need to introduce the map $\delta : \mathbb{R}^n \to \mathbb{R}$ defined by

$$\delta_i(x) = 1 - 2x_i + \|x\|^2.$$

We will also need the map $\gamma_i(\epsilon,x) : \mathbb{R}^{1,n} \to \mathbb{R}^n$ defined by

$$\gamma_i(\epsilon,x) = \delta_i(\epsilon x)^{-1}(x - \epsilon\|x\|^2e_i)$$

and the map $\kappa_i : \mathbb{R}^{1,n} \to \mathbb{R}^n$ given by

$$\kappa_i(\epsilon,x) = \delta_i(\epsilon x)^{-1}\epsilon(\epsilon x - e_i).$$

Then,

$$\exp(\epsilon\nu_i^-).f(t,x) = n_{0,\epsilon e_i}^-.\varphi(n_{t,x}) = \varphi(n_{0,-\epsilon e_i}^-.n_{t,x}).$$

Now, in order to write our result back in terms of f we need to decompose $n_{0,-\epsilon e_i}^-.n_{t,x}$ as a product of its $N \times MAN^-$ components.

Lemma 2. *For some $m \in M$*

$$n_{0,-\epsilon e_i}^-.n_{t,x} = n_{t,\gamma_i(\epsilon,x)}mh_{1,-\log(\delta_i(\epsilon x))}n_{0,\kappa_i(\epsilon,x)}^-.$$

Proof. Since $SO(n+1,1)$ has real rank 1, $W(G,A)$ has two elements. The nontrivial element $\omega \in W(G,A)$ acts on N by $\omega^{-1}N\omega = N^-$. Then, by the uniqueness of the Bruhat decomposition, it suffices to show that

$$(n_{t,\gamma_i(\epsilon,x)})^{-1}n_{0,-\epsilon e_i}^-n_{t,x}h_{1,-\log(\delta_i(\epsilon x))}^{-1}(n_{0,\kappa_i(\epsilon,x)}^-)^{-1}$$

differs from the $(n+2)\times(n+2)$ identity matrix only in the upper left $n \times n$ block (i.e. it is in M).This is a straightforward but long matrix calculation.

□

Proposition 5. *Let $f \in I'(p,r,s)$. Then,*

$$n^-_{0,-\epsilon e_i} f(t,x) = \delta_i(\epsilon x)^{-s} f(t, \gamma_i(\epsilon, x))$$

and ν_i^- acts on $I'(p,r,s)$ as the differential operator

$$(x_i^2 - \sum_{j\neq i} x_j^2)\partial_i + \sum_{j\neq i}[2x_i x_j \partial_j] + 2xsu\partial_u.$$

Proof. The first equation follows from the previous Lemma and the second equation follows form differentiating the first. ☐

We summarize the previous propositions in the following Theorem.

Theorem 1. *The action of the group G on $I'\left(p, \frac{2}{m-1}, \frac{2}{m-1}\right)$, for $p \in \mathbb{Z}_2$, gives a globalization of the action of the local symmetry group of Equation* (1.1).

Proof. Follows from Propositions 1 to 5. ☐

References

[1] R.M. Edelstein and K.S. Govinder. On a preliminary group classification of the nonlinear heat conduction equation. *Quaestiones Mathematicae*, 31(3):225–240, 2008.

[2] Jose A. Franco. Global $\widetilde{SL(2,R)}$ representations of the schrdinger equation with singular potential. *Central European Journal of Mathematics*, 10:927–941, 2012. 10.2478/s11533-012-0040-8.

[3] Maria Luz Gandarias. Classical point symmetries of a porous medium equation. *Journal of Physics A: Mathematical and General*, 29(3):607, 1996.

[4] Sepanski M. R. Hunziker, M. and R. J. Stanke. The minimal representation of the conformal group and classical solutions to the wave equation. *J. Lie Theory*, 22(2):301–360, 2012.

[5] Anthony W. Knapp. *Lie groups beyond an introduction*, volume 140 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA, 1996.

[6] Peter J. Olver. *Applications of Lie groups to differential equations*, volume 107 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1993.

[7] Mark R. Sepanski. Global actions of Lie symmetries for the nonlinear heat equation. *J. Math. Anal. Appl.*, 360(1):35–46, 2009.

[8] Mark R. Sepanski and Ronald J. Stanke. Global Lie symmetries of the heat and Schrödinger equation. *J. Lie Theory*, 20(3):543–580, 2010.

[9] Z. Wu. *Nonlinear Diffusion Equations*. World Scientific, 2001.

Department of Mathematics and Statistics, University of North Florida, 1 UNF Drive, Jacksonville, FL 32224

E-mail address: `jose.franco@unf.edu`